

Vesicular Ca^{2+} mediates granule motion and exocytosis

Ricardo Borges*, Natalia Domínguez, Judith Estévez-Herrera, Daniel Pereda, José David Machado

Unidad de Farmacología, Facultad de Medicina, Universidad de La Laguna, Tenerife, Spain

ARTICLE INFO

Article history:

Received 29 October 2011

Received in revised form 5 December 2011

Accepted 8 December 2011

Available online 4 January 2012

Keywords:

Amperometry

Bafilomycin

Calcium

Exocytosis

Secretion

TIRFM

ABSTRACT

Secretory vesicles of chromaffin cells are acidic organelles that maintain an increasing pH gradient towards the cytosol (5.5 vs. 7.3) that is mediated by V-ATPase activity. This gradient is primarily responsible for the accumulation of large concentrations of amines and Ca^{2+} , although the mechanisms mediating Ca^{2+} uptake and release from granules, and the physiological relevance of these processes, remain unclear. The presence of a vesicular matrix appears to create a bi-compartmentalised medium in which the major fractions of solutes, including catecholamines, nucleotides and Ca^{2+} , are strongly associated with vesicle proteins, particularly chromogranins. This association appears to be favoured at acidic pH values. It has been demonstrated that disrupting the pH gradient of secretory vesicles reduces their rate of exocytosis and promotes the leakage of vesicular amines and Ca^{2+} , dramatically increasing the movement of secretory vesicles and triggering exocytosis. In this short review, we will discuss the data available that highlights the importance of pH in regulating the association between chromogranins, vesicular amines and Ca^{2+} . We will also address the potential role of vesicular Ca^{2+} in two major processes in secretory cells, vesicle movement and exocytosis.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Secretory cells are characterized by the presence of secretory storage vesicles that release neurotransmitters and hormones. Adrenal chromaffin cells are commonly used to study the basic mechanisms of neurotransmitter secretion, which involves vesicle sorting, movement and exocytosis. As large concentrations of Ca^{2+} are stored in secretory vesicles, considerable effort has focused on studying the physiological role(s) of vesicular Ca^{2+} in vesicle movement and exocytosis [1–3]. The currently available experimental evidence indicates that the vesicular cocktail constitutes a source of Ca^{2+} that may be involved in regulating vesicle movement and exocytosis. Since the majority of this data is derived from experiments performed in chromaffin cells, we will first describe the most representative organelle of these cells, the chromaffin granule.

1.1. Secretory granules

Secretory granules from chromaffin cells are large, dense, core vesicles (LDCV) similar to those found in many other neuroendocrine cells and sympathetic neurons [4]. Chromaffin granules concentrate transmitters in a highly efficient manner, accumulating catecholamines (500–1000 mM [5,6]) and other soluble components such as ATP (125–300 mM [7]), ascorbate (10–30 mM [8,9]), peptides and chromogranins, thereby forming a condensed protein matrix (~180 mg/mL [10]). Furthermore, secretory granules concentrate H^+ to create an acidic medium that facilitates the accumulation of high concentrations of Ca^{2+} . The mechanisms that mediate the concentration of these elements, despite the large osmotic forces involved, have intrigued researchers for decades.

Chromaffin granules maintain a pH gradient across their membranes of about 2 orders of magnitude (from ≈ 5.5 on the inside to ≈ 7.3 in the cytosol). This gradient is maintained by the activity of a specific vesicular H^+ -ATPase (V-ATPase; [11,12]), a tightly regulated H^+ transporter present in the membranes of almost all known secretory vesicles [13,14]. This vesicular H^+ gradient acts as an antiporter to accumulate catecholamines through the vesicular monoamine transporter VMAT2 [15], and Ca^{2+} through the $\text{H}^+/\text{Ca}^{2+}$ antiporter, although most vesicular Ca^{2+} accumulation appears to occur via a SERCA-type Ca^{2+} ATPase [16,17]. The vesicular matrix composed of the distinct proteins and components of the vesicular cocktail appears to play a crucial role in chelation and in reducing the osmotic forces [18], thereby permitting solutes to accumulate at high concentrations [19]. Therefore, most of the intravesicular

Abbreviations: V-ATPase, vesicular H^+ -ATPase; VMAT2, vesicular monoamine transporter; LDCV, large dense core vesicles; Cgs, chromogranins; CgA, chromogranin A; CgB, chromogranin B; SgII, secretogranin type II; SERCA, sarcoplasmic reticulum calcium ATPase; CICR, Ca^{2+} -induced Ca^{2+} release.

* Corresponding author at: Unidad de Farmacología, Facultad de Medicina, Universidad de La Laguna, E-38071-La Laguna, Tenerife, Spain. Tel.: +34 922 319346; fax: +34 922 655995.

E-mail address: rborges@ull.es (R. Borges).

solutes are not free but rather, they are associated with other components of the vesicular matrix, the main protein components of which are chromogranins with a pK_a of about 5.5 [18,20].

1.2. Chromogranins are the main protein components of LDCV

Chromogranin A was discovered in the mid sixties [21] as the first of a series of acidic proteins generically known as granins, of which there are currently 9 examples [22,23]. In chromaffin secretory vesicles, only chromogranins A (CgA) and B (CgB) are really relevant, and to a lesser extent secretogranin II (SgII). The physiological role of Cgs as sources of biologically active peptides has been heavily investigated (see [22]), although granins can also promote granule biogenesis [24,25]. However, in non-secretory cells exogenous granins appear to accumulate in dense compartments that resemble secretory granules [24–27]. Conversely, recent studies using Cg knockout mice have shown that secretory granules persist and secretory responses are maintained in the absence of CgA [19], CgB [28] or both [29]. However, the capacity of secretory granules to accumulate solutes is significantly affected by the absence of Cgs. Although the role of Cgs in the accumulation of other solutes such as Ca^{2+} , H^+ or ATP is still unknown, the ability of Cgs to efficiently bind Ca^{2+} and H^+ suggests that the accumulation of other solutes is also impaired in the absence of Cgs.

As the association of Cgs with other solutes is also pH-dependent [18], it is plausible that intravesicular pH regulates the ability of chromogranin A to form aggregates [30], and that regulating vesicular pH plays an important role in the dynamics of vesicular Ca^{2+} and catechols [16,31,32]. The presence of a functional IP_3 receptor directly coupled to Cgs has been described, which may be involved in the physiological release of Ca^{2+} from vesicles [33]. The association of vesicular membrane-bound IP_3 -R to CgA [34] and CgB [35] is strongly dependent on pH.

1.3. Bi-compartmental storage of catecholamines and Ca^{2+}

Intravesicular Ca^{2+} was first implicated in the exocytotic process in 1967 [36], although this hypothesis is yet to be fully accepted by the scientific community. The endoplasmic reticulum has classically been considered as the main internal source of Ca^{2+} , largely because the mobilization of Ca^{2+} from intracellular stores by $InsP_3$ was first described in this organelle. More recent studies have demonstrated the involvement of other cell structures in the uptake, release and cytosolic redistribution of Ca^{2+} , including the mitochondria, nucleus and Golgi apparatus [37–39]. In this respect, secretory vesicles have received modest attention and they are frequently considered little more than a non-functional Ca^{2+} sink. The main argument for this, although with little experimental support, is that vesicular Ca^{2+} is sequestered into the vesicular matrix where it undergoes little turnover. However, recent findings and a reinterpretation of classical data appears to contradict this assumption for several reasons:

- i) Secretory granules are the most abundant organelle in chromaffin cells, as well as in many other cell types including pancreatic β -cells, mast cells and lactotrophs. Approximately 30% of the chromaffin cell volume is occupied by about 20,000 granules [40], while β -cells are estimated to contain $\approx 10,000$ insulin granules that occupy 10–20% of their total volume [41].
- ii) Chromaffin granules contain far more Ca^{2+} than any other organelle, accounting for about 60% of total Ca^{2+} in chromaffin cells [42,43]. Moreover, calcium is highly concentrated in secretory vesicles. Experiments targeting aequorins to the inside of secretory vesicles have confirmed directly that Ca^{2+} is distributed in two fractions: chelated Ca^{2+} at an estimated concentration of about 40 mM [43]; and the free Ca^{2+} fraction

at a concentration of about 50–100 μM [16,43,44]. The free fraction is equilibrated with bound Ca^{2+} , facilitating rapid recovery after acute depletion.

- iii) Vesicular Ca^{2+} is the closest source of the cation for granule movement and exocytosis.

Despite these observations, and the crucial role of this cation in processes like vesicle movement and exocytosis, the functional role(s) of vesicular Ca^{2+} has received little attention.

1.4. Mobilization of vesicular Ca^{2+}

Disruption of the pH gradient using protonophores [45] or weak bases [1,2,46] has been used to induce the alkalization of granules, resulting in the release of Ca^{2+} and catecholamines into the cytosol [46] that can promote acceleration of granule motion and its exocytosis [47]. This effect is shared by clinically used drugs such as the hypotensive agent hydralazine [48], amphetamines [49] and β -adrenergic blockers [50].

Stimuli such as histamine, caffeine and depolarization can mobilize the free Ca^{2+} fraction in vesicles [16,44]. Targeted aequorin studies suggest that intravesicular Ca^{2+} kinetics follow a bi-compartmental model, whereby a large amount of bound Ca^{2+} can rapidly replenish the free Ca^{2+} fraction after depletion induced by SERCA inhibitors (2,5-di-tert-butylhydroquinone (BHQ), cyclopiazonic acid) or pH-disrupting agents [16,44]. In addition, both $InsP_3$ - and Ca^{2+} -induced Ca^{2+} release (CICR) occur in chromaffin [16], PC12 [44] and INS-1 secretory vesicles [51]. Ca^{2+}/H^+ antiport activity has also been demonstrated in synaptic vesicles [52–55]. However, the main problem in demonstrating the participation of intravesicular Ca^{2+} in granule movement and exocytosis under physiological conditions is the difficulty in differentiating intravesicular Ca^{2+} from Ca^{2+} from other internal or external sources. All known secretagogues increase free intracellular Ca^{2+} by promoting its entry from extracellular or internal sources. Nevertheless, the vesicular alkalization observed upon the activation of several second messenger pathways also contributes to the mobilization of vesicular Ca^{2+} and catecholamines (this latter effect having recently been demonstrated using single cell amperometry: [46,47]). Nevertheless, it is plausible that the pH gradient across the vesicular membrane provides the necessary link between physiological stimuli and the regulation of Ca^{2+} and catecholamine release from secretory vesicles.

The presence of these rapid release mechanisms in secretory vesicles is frequently ignored. Conversely, data from experiments using pH-disrupting agents or drugs that act on ryanodine- or IP_3 -receptors have been interpreted as reflecting specific events associated to the endoplasmic reticulum or mitochondria, organelles of lesser importance in terms of Ca^{2+} capacity than secretory vesicles, and generally more distant. Thus, in addition to investigating the role of other known organelles, future studies using cell stimulation (via $InsP_3$ receptors, ryanodine receptors or plasma membrane Ca^{2+} channels) should take into account the induction of vesicular Ca^{2+} release. Moreover, other stimuli that activate guanylate cyclase or adenylate cyclase, which alkalize the vesicular lumen, may also mimic these mechanisms. Given the poor diffusion of Ca^{2+} through the cytosol [56], it is highly plausible that vesicular Ca^{2+} plays an important physiological role in the granule's approach to the membrane [3,57,58] and subsequent exocytosis. To examine this proposal we studied the effects of bafilomycin A1, a potent and highly specific inhibitor of the H^+ -ATPase, on vesicle alkalization and the release of vesicular Ca^{2+} into the cytosol. This Ca^{2+} release increases the lateral movement of chromaffin granules and triggers exocytosis. Although bafilomycin is not a physiological stimulus, these results reveal a novel mechanism for vesicular Ca^{2+} release controlled by the pH gradient.




Fig. 1. Mechanism of Ca^{2+} (and catecholamine, CA) turnover in chromaffin secretory organelles. The relative sizes of the granule matrix (1) and the free compartment (2) have been altered for clarity. H^+ is pumped towards the vesicle lumen by an ATP dependent pump (V-ATPase, 3). Protons maintain the pH and the potential gradients with the help of Cl^- channels, Cl^- ions acting as counter ions (4) to maintain a membrane potential at approximately -80 mV . Catecholamines (5) and Ca^{2+} (6) use the H^+ gradient as an antiporter to accumulate in vesicles. Both antiporters can also work in reverse mode. IP_3 receptors (7) mediate Ca^{2+} release in response to intracellular IP_3 , whereas CICR (8) enhances Ca^{2+} signaling via ryanodine- and caffeine-modulated Ca^{2+} release. The SERCA (9), which to date has not been described in chromaffin granules, acts as a Ca^{2+} pump that can be blocked by thapsigargin. In these studies the luminal terminal of VAMP (10) (synaptobrevin) has been modified to accommodate a Ca^{2+} (low Ca^{2+} -affinity aequorin) or pH (EGFP) sensor.

Adapted from Ref. [3].

Further studies will be necessary to determine the physiological relevance of Ca^{2+} release from secretory vesicles.

In summary, when our recent findings are considered in the light of those from other laboratories, it appears that: (i) secretory vesicles from PC12 [44] and chromaffin cells [16] accumulate Ca^{2+} in two distinct and exchangeable forms, free ($\approx 50\text{--}100\ \mu\text{M}$) and bound Ca^{2+} ($\approx 40\ \text{mM}$); (ii) vesicular pH is closely associated with the modulation of the kinetics and quantal characteristics of catecholamine exocytosis [46]; (iii) secretory granules possess mechanisms for fast uptake and release of Ca^{2+} ; (iv) Ca^{2+} release from granules can influence granule movement and exocytosis [47]. The main mechanisms underlying vesicular Ca^{2+} turnover are depicted in Fig. 1.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Acknowledgments

ND is the recipient of an FPU fellowship, while JE and DP are recipients of FPI fellowships from the Spanish Ministry of Science and Innovation (MICINN). JDM holds a Ramon y Cajal contract from the MICINN. This work was supported by MICINN (BFU2007-64963), Consolider (CSD2008-00005) and the Canary Islands Agency for Research, Innovation and Information Society PI2007/017. We are grateful to all past members of our research group. Discussions with Mark Sefton were greatly appreciated.

References

[1] M.L. Mundorf, S.E. Hochstetler, R.M. Wightman, Amine weak bases disrupt vesicular storage and promote exocytosis in chromaffin cells, *J. Neurochem.* 73 (1999) 2397–2405.

[2] M.L. Mundorf, K.P. Troyer, S.E. Hochstetler, J.A. Near, R.M. Wightman, Vesicular Ca^{2+} participates in the catalysis of exocytosis, *J. Biol. Chem.* 275 (2000) 9136–9142.

[3] J.D. Machado, M. Camacho, J. Alvarez, R. Borges, On the role of intravesicular calcium in the motion and exocytosis of secretory organelles, *Commun. Integr. Biol.* 2 (2009) 71–73.

[4] H. Winkler, The adrenal chromaffin granule: a model for large dense core vesicles of endocrine and nervous tissue, *J. Anat.* 183 (Pt 2) (1993) 237–252.

[5] J.A. Jankowski, T.J. Schroeder, E.L. Ciolkowski, R.M. Wightman, Temporal characteristics of quantal secretion of catecholamines from adrenal-medullary cells, *J. Biol. Chem.* 268 (1993) 14694–14700.

[6] A. Albillos, G. Dernick, H. Horstmann, W. Almers, G. Alvarez de Toledo, M. Lindau, The exocytotic event in chromaffin cells revealed by patch amperometry, *Nature* 389 (1997) 509–512.

[7] A. Weber, H. Winkler, Specificity and mechanism of nucleotide uptake by adrenal chromaffin granules, *Neuroscience* 6 (1981) 2269–2276.

[8] O. Terland, T. Flatmark, Ascorbate as a natural constituent of chromaffin granules from the bovine adrenal medulla, *FEBS Lett.* 59 (1975) 52–56.

[9] F.S. Menniti, J. Knoth, D.S. Peterson, E.J. Diliberto Jr., The in situ kinetics of dopamine beta-hydroxylase in bovine adrenomedullary chromaffin cells. Intravesicular compartmentation reduces apparent affinity for the cofactor ascorbate, *J. Biol. Chem.* 262 (1987) 7651–7657.

[10] J.H. Phillips, Passive ion permeability of the chromaffin-granule membrane, *Biochem. J.* 168 (1977) 289–297.

[11] N. Nelson, Structure and pharmacology of the proton-ATPases, *Trends Pharmacol. Sci.* 12 (1991) 71–75.

[12] S. Saroussi, N. Nelson, The little we know on the structure and machinery of V-ATPase, *J. Exp. Biol.* 212 (2009) 1604–1610.

[13] S.L. Gluck, The structure and biochemistry of the vacuolar H^+ ATPase in proximal and distal urinary acidification, *J. Bioenerg. Biomembr.* 24 (1992) 351–359.

[14] M. Toei, R. Saum, M. Forgac, Regulation and isoform function of the V-ATPases, *Biochemistry* 49 (2010) 4715–4723.

[15] L. Johannes, P.M. Lledo, M. Roa, J.D. Vincent, J.P. Henry, F. Darchen, The GTPase Rab3a negatively controls calcium-dependent exocytosis in neuroendocrine cells, *EMBO J.* 13 (1994) 2029–2037.

[16] J. Santodomingo, L. Vay, M. Camacho, E. Hernandez-Sanmiguel, R.I. Fonteriz, C.D. Lobaton, M. Montero, A. Moreno, J. Alvarez, Calcium dynamics in bovine adrenal medulla chromaffin cell secretory granules, *Eur. J. Neurosci.* 28 (2008) 1265–1274.

[17] J.R. Haigh, J.H. Phillips, Indirect coupling of calcium transport in chromaffin granule ghosts to the proton pump, *Neuroreport* 4 (1993) 571–574.

[18] K.B. Helle, R.K. Reed, K.E. Pihl, G. Serck-Hanssen, Osmotic properties of the chromogranins and relation to osmotic pressure in catecholamine storage granules, *Acta Physiol. Scand.* 123 (1985) 21–33.

[19] M.S. Montesinos, J.D. Machado, M. Camacho, J. Diaz, Y.G. Morales, D. Alvarez de la Rosa, E. Carmona, A. Castaneya, O.H. Viveros, D.T. O'Connor, S.K. Mahata, R. Borges, The crucial role of chromogranins in storage and exocytosis revealed using chromaffin cells from chromogranin A null mouse, *J. Neurosci.* 28 (2008) 3350–3358.

[20] J.S. Videen, M.S. Mezger, Y.M. Chang, D.T. O'Connor, Calcium and catecholamine interactions with adrenal chromogranins. Comparison of driving forces in binding and aggregation, *J. Biol. Chem.* 267 (1992) 3066–3073.

[21] H. Blaschko, R.S. Comline, F.H. Schneider, M. Silver, A.D. Smith, Secretion of a chromaffin granule protein, chromogranin, from the adrenal gland after splanchnic stimulation, *Nature* 215 (1967) 58–59.

[22] L. Taupenot, K.L. Harper, D.T. O'Connor, The chromogranin-secretogranin family, *N. Engl. J. Med.* 348 (2003) 1134–1149.

[23] R. Borges, J. Diaz-Vera, N. Dominguez, M.R. Arnau, J.D. Machado, Chromogranins as regulators of exocytosis, *J. Neurochem.* 114 (2010) 335–343.

[24] T. Kim, J.H. Tao-Cheng, L.E. Eiden, Y.P. Loh, Chromogranin A, an “on/off” switch controlling dense-core secretory granule biogenesis, *Cell* 106 (2001) 499–509.

[25] Y.H. Huh, S.H. Jeon, S.H. Yoo, Chromogranin B-induced secretory granule biogenesis: comparison with the similar role of chromogranin A, *J. Biol. Chem.* 278 (2003) 40581–40589.

[26] N. Beuret, H. Stettler, A. Renold, J. Rutishauser, M. Spiess, Expression of regulated secretory proteins is sufficient to generate granule-like structures in constitutively secreting cells, *J. Biol. Chem.* 279 (2004) 20242–20249.

[27] M.L. Malosio, T. Giordano, A. Laslop, J. Meldolesi, Dense-core granules: a specific hallmark of the neuronal/neurosecretory cell phenotype, *J. Cell Sci.* 117 (2004) 743–749.

[28] J. Diaz-Vera, Y.G. Morales, J.R. Hernandez-Fernaund, M. Camacho, M.S. Montesinos, F. Calegari, W.B. Huttner, R. Borges, J.D. Machado, Chromogranin B gene ablation reduces the catecholamine cargo and decelerates exocytosis in chromaffin secretory vesicles, *J. Neurosci.* 30 (2010) 950–957.

[29] J. Diaz-Vera, M. Camacho, J. Machado, N. Dominguez, M. Montesinos, J. Hernandez-Fernaund, R. Lujan, R. Borges, Chromogranins A and B are key proteins in amine accumulation but the catecholamine secretory pathway is conserved without them, *FASEB J.* (2012) 26.

[30] L. Taupenot, K.L. Harper, D.T. O'Connor, Role of H^+ -ATPase-mediated acidification in sorting and release of the regulated secretory protein chromogranin A: evidence for a vesiculogenic function, *J. Biol. Chem.* 280 (2005) 3885–3897.

[31] S.H. Yoo, J.P. Albanesi, High capacity, low affinity Ca^{2+} binding of chromogranin A. Relationship between the pH-induced conformational change and Ca^{2+} binding property, *J. Biol. Chem.* 266 (1991) 7740–7745.

- [32] N.R. Mahapatra, M. Mahata, P.P. Hazra, P.M. McDonough, D.T. O'Connor, S.K. Mahata, A dynamic pool of calcium in catecholamine storage vesicles. Exploration in living cells by a novel vesicle-targeted chromogranin A-aequorin chimeric photoprotein, *J. Biol. Chem.* 279 (2004) 51107–51121.
- [33] S.H. Yoo, S.Y. Chu, K.D. Kim, Y.H. Huh, Presence of secretogranin II and high-capacity, low-affinity Ca^{2+} storage role in nucleoplasmic Ca^{2+} store vesicles, *Biochemistry* 46 (2007) 14663–14671.
- [34] S.H. Yoo, pH-dependent association of chromogranin A with secretory vesicle membrane and a putative membrane binding region of chromogranin A, *Biochemistry* 32 (1993) 8213–8219.
- [35] S.H. Yoo, pH-dependent binding of chromogranin B and secretory vesicle matrix proteins to the vesicle membrane, *Biochim. Biophys. Acta* 1179 (1993) 239–246.
- [36] J.L. Borowitz, Calcium binding by subcellular fractions of bovine adrenal medulla, *J. Cell. Physiol.* 69 (1967) 311–319.
- [37] C. Villalobos, L. Nunez, M. Montero, A.G. Garcia, M.T. Alonso, P. Chamero, J. Alvarez, J. Garcia-Sancho, Redistribution of Ca^{2+} among cytosol and organelle during stimulation of bovine chromaffin cells, *FASEB J.* 16 (2002) 343–353.
- [38] M. Montero, M.T. Alonso, E. Carnicero, I. Cuchillo-Ibanez, A. Albillos, A.G. Garcia, J. Garcia-Sancho, J. Alvarez, Chromaffin-cell stimulation triggers fast millimolar mitochondrial Ca^{2+} transients that modulate secretion, *Nat. Cell Biol.* 2 (2000) 57–61.
- [39] A.G. Garcia, A.M. Garcia-De-Diego, L. Gandia, R. Borges, J. Garcia-Sancho, Calcium signaling and exocytosis in adrenal chromaffin cells, *Physiol. Rev.* 86 (2006) 1093–1131.
- [40] H. Plattner, A.R. Artalejo, E. Neher, Ultrastructural organization of bovine chromaffin cell cortex-analysis by cryofixation and morphometry of aspects pertinent to exocytosis, *J. Cell Biol.* 139 (1997) 1709–1717.
- [41] P.M. Dean, Ultrastructural morphometry of the pancreatic-cell, *Diabetologia* 9 (1973) 115–119.
- [42] J.R. Haigh, R. Parris, J.H. Phillips, Free concentrations of sodium, potassium and calcium in chromaffin granules, *Biochem. J.* 259 (1989) 485–491.
- [43] D. Bulenda, M. Gratzl, Matrix free Ca^{2+} in isolated chromaffin vesicles, *Biochemistry* 24 (1985) 7760–7765.
- [44] A. Moreno, C.D. Lobaton, J. Santodomingo, L. Vay, E. Hernandez-SanMiguel, R. Rizzuto, M. Montero, J. Alvarez, Calcium dynamics in catecholamine-containing secretory vesicles, *Cell Calcium* 37 (2005) 555–564.
- [45] C.L. Haynes, L.A. Buhler, R.M. Wightman, Vesicular $\text{Ca}(2+)$ -induced secretion promoted by intracellular pH-gradient disruption, *Biophys. Chem.* 123 (2006) 20–24.
- [46] M. Camacho, J.D. Machado, M.S. Montesinos, M. Criado, R. Borges, Intracellular pH rapidly modulates exocytosis in adrenal chromaffin cells, *J. Neurochem.* 96 (2006) 324–334.
- [47] M. Camacho, J.D. Machado, J. Alvarez, R. Borges, Intravesicular calcium release mediates the motion and exocytosis of secretory organelles: a study with adrenal chromaffin cells, *J. Biol. Chem.* 283 (2008) 22383–22389.
- [48] J.D. Machado, J.F. Gomez, G. Betancor, M. Camacho, M.A. Brioso, R. Borges, Hydralazine reduces the quantal size of secretory events by displacement of catecholamines from adrenomedullary chromaffin secretory vesicles, *Circ. Res.* 91 (2002) 830–836.
- [49] D. Sulzer, T.K. Chen, Y.Y. Lau, H. Kristensen, S. Rayport, A. Ewing, Amphetamine redistributes dopamine from synaptic vesicles to the cytosol and promotes reverse transport, *J. Neurosci.* 15 (1995) 4102–4108.
- [50] M.S. Montesinos, M. Camacho, J.D. Machado, O.H. Viveros, B. Beltran, R. Borges, The quantal secretion of catecholamines is impaired by the accumulation of beta-adrenoceptor antagonists into chromaffin cell vesicles, *Br. J. Pharmacol.* 159 (2010) 1548–1556.
- [51] J. SantoDomingo, R.I. Fonteriz, C.D. Lobaton, M. Montero, A. Moreno, J. Alvarez, Ca^{2+} dynamics in the secretory vesicles of neurosecretory PC12 and INS1 cells, *Cell. Mol. Neurobiol.* 30 (2010) 1267–1274.
- [52] P.P. Goncalves, S.M. Meireles, C. Gravato, M.G. Vale, Ca^{2+} - H^{+} antiport activity in synaptic vesicles isolated from sheep brain cortex, *Neurosci. Lett.* 247 (1998) 87–90.
- [53] P.P. Goncalves, S.M. Meireles, P. Neves, M.G. Vale, Synaptic vesicle $\text{Ca}^{2+}/\text{H}^{+}$ antiport: dependence on the proton electrochemical gradient, *Brain Res. Mol. Brain Res.* 71 (1999) 178–184.
- [54] P.P. Goncalves, S.M. Meireles, P. Neves, M.G. Vale, Ionic selectivity of the $\text{Ca}^{2+}/\text{H}^{+}$ antiport in synaptic vesicles of sheep brain cortex, *Brain Res. Mol. Brain Res.* 67 (1999) 283–291.
- [55] P.P. Goncalves, S.M. Meireles, P. Neves, M.G. Vale, Distinction between $\text{Ca}(2+)$ pump and $\text{Ca}(2+)/\text{H}(+)$ antiport activities in synaptic vesicles of sheep brain cortex, *Neurochem. Int.* 37 (2000) 387–396.
- [56] F. Sala, A. Hernandez-Cruz, Calcium diffusion modeling in a spherical neuron. Relevance of buffering properties, *Biophys. J.* 57 (1990) 313–324.
- [57] L. von Ruden, E. Neher, A Ca-dependent early step in the release of catecholamines from adrenal chromaffin cells, *Science* 262 (1993) 1061–1065.
- [58] M.W. Allersma, M.A. Bittner, D. Axelrod, R.W. Holz, Motion matters: secretory granule motion adjacent to the plasma membrane and exocytosis, *Mol. Biol. Cell* 17 (2006) 2424–2438.